## Rehabilitation Services, Inc.

42 Lloyd Avenue, Suite A, Malvern, PA 19355-3000 610 644 0136

## INFORMED CONSENT for ASSESSMENT AND TREATMENT OF THE PELVIC FLOOR

Internal examination of the pelvic floor muscles is consistent with physical therapy practice. It complies with national physical therapy policies requiring the performance of test and measurements of neuromuscular function as an aid to the evaluation and treatment of a specific medical condition.

-this statement was adopted by the executive committee of the Section on Obstetrics and Gynecology of the American Physical therapy Association

-San Antonio, Texas.....February 1993

I understand that it may be beneficial for my therapist to perform soft tissue assessment and treatment of the pelvic floor. Palpation of this area is most direct and accessible if done via the vagina and/or rectum. Pelvic floor dysfunctions include pelvic pain, urinary or fecal incontinence, dyspareunia (pain with intercourse), pain from episiotomy or scarring, vulvodynia, vestibulitis, or other similar conditions. Restrictions in this area may also be contributing to symptoms in other areas of the body.

I understand that the benefits of this procedure will be explained to me. I understand that, if I am uncomfortable with participating in this treatment procedure AT ANY TIME, I will inform the therapist and the procedure will be discontinued and alternatives will be discussed with me.

This direct pelvic floor release procedure utilizes Myofascial Release principles entailing the relaxation and lengthening of muscles, fascia and other soft tissue structures within the areas of the pelvic floor, sacrum, coccyx, and the sacroiliac, hip and pubic joints. The procedure also requires pressure and/or distraction directly to the coccyx bone. This technique is an accepted physical therapy technique, as indicated above. Our experience has demonstrated that this direct pelvic floor release is helpful, often facilitating consistent therapeutic results. As with any area of the body, most people require a series of these specific treatments. This is determined by evaluation and treatment findings.

Myofascial Release Treatment Center.	e above procedure being performed by	y the merapisis at the
Patient's Printed Name	Date	
Patient's Signature		
Witness or Therapist's Signature		

\*\*\*If you are pregnant, have infections of any kind, have vaginal dryness, are less than 6 weeks post partum or post surgery, have severe pelvic pain, using any IUD, sensitivity to KY jelly, vaginal creams or latex, please inform the therapist prior to this procedure.